

Recent advances in crop transformation technologies

Received: 28 February 2022

Accepted: 30 October 2022

Published online: 15 December 2022

 Check for updates

Zongliang Chen ¹, Juan M. Debernardi ^{2,3}, Jorge Dubcovsky ^{2,3}
& Andrea Gallavotti ^{1,4} 

Agriculture is experiencing a technological inflection point in its history, while also facing unprecedented challenges posed by human population growth and global climate changes. Key advancements in precise genome editing and new methods for rapid generation of bioengineered crops promise to both revolutionize the speed and breadth of breeding programmes and increase our ability to feed and sustain human population growth. Although genome editing enables targeted and specific modifications of DNA sequences, several existing barriers prevent the widespread adoption of editing technologies for basic and applied research in established and emerging crop species. Inefficient methods for the transformation and regeneration of recalcitrant species and the genotype dependency of the transformation process remain major hurdles. These limitations are frequent in monocotyledonous crops, which alone provide most of the calories consumed by human populations. Somatic embryogenesis and de novo induction of meristems – pluripotent groups of stem cells responsible for plant developmental plasticity – are essential strategies to quickly generate transformed plants. Here we review recent discoveries that are rapidly advancing nuclear transformation technologies and promise to overcome the obstacles that have so far impeded the widespread adoption of genome editing in crop species.

The efficient production of transgenic plants relies on two key steps: transformation (the transfer and expression of transgenes into host cells) and regeneration (the ability to form a fertile plant from a transformed cell). For many species, transformation and regeneration are the bottlenecks to obtaining transgenic plants. Traditionally, callus induction from tissue explants has been the main avenue to produce transformed plants, either by providing undifferentiated cells for direct transformation or as a way to regenerate full plants from a few transformed cells (Fig. 1). However, callus formation is a lengthy procedure known to introduce genomic and epigenomic changes, often with unintended consequences^{1,2}. Recent progress in both transformation and regeneration promises to quickly advance our ability to manipulate crop genomes. This includes the use of different morphogenic factors,

genes that are involved in somatic embryogenesis or meristem development, to trigger reprogramming and pluripotency of a subset of somatic cells to eventually produce transformed plants³.

Methods for plant transformation

The efficient delivery of gene-modification components into plant cells is a crucial first step of plant transformation and genome editing. Two major gene-delivery methodologies have been established in higher plants since the 1980s: direct gene transfer and *Agrobacterium*-mediated transformation⁴ (Fig. 1). Direct gene transfer methods are good strategies to overcome possible competency barriers to transformation, and traditionally, microprojectile bombardment (also known as biolistics) has been used to facilitate the delivery of

¹Waksman Institute of Microbiology, Rutgers University, Piscataway, NJ, USA. ²Department of Plant Sciences, University of California Davis, Davis, CA, USA. ³Howard Hughes Medical Institute, Chevy Chase, MD, USA. ⁴Department of Plant Biology, Rutgers University, New Brunswick, NJ, USA.

 e-mail: agallavotti@waksman.rutgers.edu




Fig. 1 | Plant transformation and regeneration strategies. **a**, *Agrobacterium*-mediated transformation strategies include callus induction followed by organogenesis (traditional transformation), somatic embryogenesis and de novo shoot regeneration in planta promoted by morphogenic genes (fast

transformation). **b**, Particle bombardment for nuclear plant transformation. **c**, Various nanoparticles and virus-based RNA delivery systems for plant transformation. Created with BioRender.com.

transgenes in many species. Plants obtained from particle bombardments can contain multiple integration events and random rearrangements of the integrated copies with unpredictable effects, complicating the downstream analysis of the transgenic plants generated⁵. However, these phenomena are not unique to biolistic methods; they are sometimes observed in *Agrobacterium*-mediated transformations as well^{6–10}. Recently, biolistic delivery has been used to directly provide in vitro transcripts or ribonucleoprotein complexes of CRISPR–Cas9 to regenerable plant tissues of a wider range of genotypes and species, obviating genome integration effects^{11–14}.

Agrobacterium-mediated transformation is a cost-effective, efficient gene-delivery system, capable of transferring large DNA fragments into plant chromosomes¹⁵, and remains the top choice for plant transformation. However, only a small range of plant host genotypes are competent for *Agrobacterium* infection, and in numerous monocot crops, ineffective *Agrobacterium* infection hinders its widespread application. To improve transformation efficiency, substantial efforts have been made to optimize virulence (*vir*) gene expression of *Agrobacterium* strains. Progress in our understanding of the mechanisms used by

Agrobacterium to modulate plant host defence responses for effective infection and of the role of phytohormones in the early stages of plant–*Agrobacterium* interaction is summarized in a recent review¹⁶. The process of T-DNA transfer starts with a two-component system containing the hybrid histidine kinase VirA and the response regulator VirG. The VirA protein is activated by phenolic compounds (such as acetosyringone, a common ingredient in transformation media) and triggers the phosphorylation of VirG, which induces the expression of other *vir* genes in the Tumor-inducing (Ti) plasmid^{16,17}. The *vir* gene expression can be repressed by phytohormones from the host side at the early stage of plant–*Agrobacterium* interaction. In particular, salicylic acid (SA) prevents the induction of *vir* genes by attenuating the kinase function of the VirA protein^{16,18}. Consequently, plants that overproduce SA are recalcitrant to *Agrobacterium* infections, whereas those with defective SA biosynthesis are easier to transform^{18,19}. The negative effect of SA on the VirA/G two-component regulatory system can be alleviated by the addition of acetosyringone¹⁸. Different *Agrobacterium* strains containing binary, super-binary and recently emerged ternary vectors have been developed to fully exploit the

power of the *vir* genes¹⁶. Specifically, improved ternary vector systems introduce a third helper plasmid containing extra *vir* genes and have been shown to enhance transformation efficiency in recalcitrant maize lines^{20,21}. These strains, combined with a fast-transformation protocol, increased transformation efficiency in the commonly used maize inbred line B104 (ref. 22). Recently, engineered *Agrobacterium* strains expressing a type III secretion system to deliver *Pseudomonas* effectors that repress host defence responses resulted in increased transformation efficiency in wheat, alfalfa and switchgrass²³.

Nanoparticle²⁴ and virus-based RNA delivery systems can bypass tissue culture and are attracting increasing attention for efficient genome editing. Several nanomaterials, such as silica, metal, polymeric and magnetic nanoparticles as well as carbon nanotubes, have been investigated for their potential role in gene delivery for plant transformation. Although recently questioned²⁵, a novel transformation methodology called pollen magnetofection was reported to allow the stable integration of transgenes in the nuclear genome of several dicot species²⁶. In this technology, recombinant DNA is attached on the positively charged surface of magnetic nanoparticles that can penetrate pollen grains in a magnetic field; subsequent fertilizations are conducted to complete the transformation process^{26,27}.

Virus-based RNA delivery systems have the additional advantage of bypassing transgene integration, thereby avoiding possible disruptive effects on gene function. RNA-virus-based vector systems have been developed to express guide RNA arrays for multiplex genome editing in several species. In tobacco, a rhabdoviral system was developed to deliver all CRISPR–Cas9 components during infections; however, a lack of editing in the germline required a tissue-culture step to obtain edited progenies²⁸. An alternative approach was pursued in both wheat and maize in which the viral system relied on infections of previously obtained Cas9-positive plants, and these systems generated edited plants with high efficiency in both species^{29,30}. These virus-based systems are still limited by host range and physical constraints but are nonetheless very promising approaches for genome editing of crop species and for breeding purposes.

Molecular mechanisms of plant regeneration

Although the remarkable capacity to regenerate organs and entirely new individuals is considered a signature feature of plants³¹, many crop species are incapable of naturally regenerating whole plantlets upon loss or injury of body parts³². Nevertheless, *in vitro* cultivation of various types of explants on enriched media supplemented with specific phytohormones enables many plant species to regenerate plantlets (Fig. 1). The regeneration process typically involves a callus-formation step followed by organ differentiation, direct embryogenesis from somatic tissues (also known as somatic embryogenesis) or *de novo* formation of meristems. The capacity to regenerate meristems or whole plantlets from non-zygotic cells is the fundamental basis for agricultural and horticultural biotechnology, and it allows both clonal propagation and the production of stable transgenic material for breeding. Regardless of the substantial technological progress that has been made over the past four decades, there are still many economically important crops that are recalcitrant to *in vitro* regeneration from somatic tissue. Understanding the molecular mechanisms governing plant regeneration is therefore particularly critical for accelerating plant biotechnology.

Auxin and cytokinin are key hormones for shoot and root apical meristem formation and *de novo* organogenesis^{31,33,34}. The fate of explants under *in vitro* cultivation follows the golden hormonal regeneration rule that has been successfully applied to *in vitro* propagation of many plant species: a high cytokinin-to-auxin ratio stimulates shoot formation, while a reverse ratio promotes root formation. Auxin is a versatile regulator and plays an essential role in embryogenesis by establishing both the shoot apical and root apical meristems³⁵. In particular, auxin biosynthetic genes, such as *TRYPTOPHAN AMINOTRANSFERASE OF ARABIDOPSIS* (*TAA1*), *TAA-RELATED 1* and *2* (*TAR1,2*), and

several flavin monooxygenase-like *YUCCA* (*YUC*) genes, regulate early embryogenesis^{36,37}. In practice, auxin (mostly the synthetic auxin-like hormone 2,4-dichlorophenoxyacetic acid (2,4-D)) typically serves as a major inducer for somatic embryogenesis from explants of different origin. During 2,4-D-induced somatic embryogenesis in *Arabidopsis*, genome-wide changes in chromatin accessibility, enhanced expression of auxin pathway genes and changes in key developmental regulators are observed in a hierarchical cascade upstream of early embryonic patterning genes^{38–40} (Fig. 2). Increasing evidence points to the importance of locally elevated endogenous auxin biosynthesis, mainly indole-3-acetic acid, as key to somatic embryogenesis^{41–43}. Although the specific mechanism by which auxin biosynthesis promotes somatic embryogenesis has not been fully resolved, auxin biosynthetic genes are direct targets of the B3 transcription factor *LEAFY COTYLEDON 2* (*LEC2*), whose inducible overexpression is sufficient to promote *Arabidopsis* somatic embryogenesis independent of tissue competence, which is normally restricted to immature zygotic embryos^{40,44}. A similar stimulating effect on somatic embryogenesis was originally reported with the overexpression of *LEC1*, another transcriptional regulator⁴⁵.

Cytokinins are adenine-derived molecules that control meristem size by directly regulating the expression levels of the homeobox transcription factor *WUSCHEL* (*WUS*), which plays a central role in meristem establishment and maintenance in most species. Extensive studies in *Arabidopsis* have shown that cytokinin biosynthesis and signalling provide positional cues for *WUS* patterning in meristems via both positive and negative feedback loops³¹. During *in vitro* shoot regeneration experiments in *Arabidopsis*, the treatment of explants with high concentrations of cytokinin removed the repressive H3K27me3 histone modification at the *WUS* locus and allowed transcriptional activators of the cytokinin signal transduction pathway to promote *WUS* transcription for *de novo* formation of shoot meristems⁴⁶. ATAC-seq and ChIP-seq profiling during callus induction and shoot regeneration in *Arabidopsis* provide a molecular timeline where auxin and cytokinin contribute to the dynamic regulation of chromatin state and the accessibility of key transcription factors involved in both promoting pluripotency acquisition during callus induction (in a high-auxin/cytokinin environment) and shoot fate determination during regeneration (in low-auxin/cytokinin conditions)⁴⁷.

Wounding is a primary signal not only for naturally occurring organ repair but also for plantlet regeneration. *In vitro* shoot regeneration experiments from root explants of *Arabidopsis* have shown that this process typically involves transcriptional activation of multiple developmental regulators, including the AP2/ERF transcription factors *ENHANCER OF SHOOT REGENERATION 1* (*ESR1*) and *WOUND INDUCED DEDIFFERENTIATION 1* to *4* (*WIND1* to *4*). *WIND1* was found to activate the expression of *ESR1* and promote shoot regeneration^{48,49}. Another AP2/ERF transcription factor, *ETHYLENE RESPONSE FACTOR115* (*ERF115*), forms a heterodimeric complex with *PHYTOCHROME A SIGNAL TRANSDUCTION1* (*PAT1*) to re-establish a stem cell niche upon root tip excision, and its high regenerative potential is correlated with the activation of *WIND1*, one of its putative targets⁵⁰. Moreover, transcriptomic analysis on *Arabidopsis* hypocotyl excisions revealed that three additional AP2/ERF transcription factors, *PLETHORA 3* (*PLT3*), *PLT5* and *PLT7*, are wound-induced and promote callus formation⁵¹. *PLT* genes lead to the induction of *YUC1* and auxin-dependent activation of cell cycle regulators during callus formation from *Arabidopsis* protoplasts, providing a mechanistic model of pluripotency acquisition in differentiated cells⁵². These three factors are key regulators of lateral root development, and the transient induction of *PLT5* or *PLT7* from various *Arabidopsis* explants (or *PLT5* in snapdragon and tomato) is sufficient to trigger *de novo* shoot formation in a hormone-independent manner^{53–55}.

Wounding is also an important signal for plant glutamate-receptor-like proteins (GLRs), which have a well-documented role in defence responses. The role of glutamate receptors in regeneration




Fig. 2 | Plant regeneration pathways. Known molecular components of plant regeneration pathways. Solid lines indicate well-established relationships, while dashed lines indicate relationships where key molecular details are unknown; arrowhead ends indicate positive regulation and perpendicular ends indicate negative regulation. Created with BioRender.com.

is mediated by SA signalling, and mutants in the SA receptor NPR1 are hyper-regenerative and partially resistant to GLR perturbations, suggesting that a higher efficiency of *Agrobacterium*-mediated transformation could be obtained by attenuating the SA response. Indeed, genetic and pharmacological inhibition of GLR activity increased the regeneration efficiency of multiple organ repair systems in *Arabidopsis* and the callus induction rate of the recalcitrant maize inbred line B73. These results suggest that the regulation of the trade-off between defence and regeneration can be harnessed to improve regeneration for agricultural purposes⁵⁶ (Fig. 2).

Somatic embryogenesis shares several developmental steps with zygotic embryogenesis, and key trigger genes of embryogenesis initiation can be manipulated to induce somatic embryogenesis for whole-plant regeneration. The AP2/ERF transcription factor BABY BOOM (BBM, also known as PLT4 in *Arabidopsis*) promotes somatic embryogenesis^{57–59} and directly regulates several auxin biosynthesis genes, including YUC genes in both monocot and dicot species. Pharmacological and genetic inactivation of endogenous YUC activity drastically reduced somatic embryogenesis via BBM induction in both rice and *Arabidopsis*, indicating that endogenous indole-3-acetic acid levels cannot be substituted by the 2,4-D present in the medium^{41,60}. A recent study suggests that this may be due to differential effects of exogenous and endogenous auxin on cell cycle progression⁵². During somatic embryogenesis, YUC-dependent auxin biosynthesis does not appear to be required for re-establishing pluripotent cells but remains essential for embryo identity and growth⁴¹. BBM also functions

directly upstream of the above-mentioned transcription factor LEC2 and is in turn upregulated by LEC2 in a reinforcing loop during somatic embryogenesis^{40,61}. In rice, the *BBM1* gene can promote parthenogenesis, the formation of embryos without fertilization, when ectopically expressed in unfertilized egg cells⁵⁸. Similarly, the recently isolated *PARTHENOGENESIS* (*PAR*) gene of apomictic dandelion, encoding a K2-2 zinc finger and EAR-domain containing protein, can trigger embryogenesis in unfertilized egg cells, and its ectopic expression in unfertilized egg cells of lettuce is sufficient to induce the formation of haploid embryos⁶². While yet to be determined, it is possible that *PAR* and other parthenogenic genes could be used to promote somatic embryogenesis for whole-plant regeneration in certain species.

The roles of other genes, such as *FUSCA 3*, *LEC1* and *2*, and *AGAMOUS-LIKE 15* (*AGL15*), in promoting somatic embryogenesis have been well described in *Arabidopsis* and other species^{3,63} (Table 1), and the expression of these morphogenic genes is believed to be regulated by BBM^{61,64}. However, whether these genes can be applied to stimulate regeneration in a broad range of species has yet to be determined.

Morphogenic factors for crop transformation

A significant breakthrough technology in plant transformation, especially for monocot species and species recalcitrant to transformation, has been the exploitation of specific morphogenic factors to reprogram somatic cells into initiating embryogenesis. This has spurred a renewed interest in exploiting specific developmental regulators for crop transformation^{3,63,65} (Table 1). These factors include BBM, WUS

Table 1 | Successful examples of utilization of morphogenic genes for transformation of crop species

Individual genes	Gene name	Transformed and regenerated crop	Relevant notes	References
AGL15	GmAGL15	Soybean	Promotes somatic embryogenesis.	94
WUS/WOX	TaWOX5	Wheat, maize, triticale, rice, barley	With ternary vector system.	70,77
	AtWUS	Coffee	Promotes somatic embryogenesis.	95
	ZmWUS2	Sorghum, maize	With ternary vector system.	68,96
	BrrWUSa	Turnip	Callus step required.	97
PLT	AtPLT5	Tomato, snapdragon	Organogenesis at wound site.	55
KN1/STM	ZmKN1	Citrus	Promotes organogenesis.	98
	BnSTM BoSTM BrSTM	Rapeseed	Promotes microspore-derived embryogenesis.	99
BBM	BnBBM	Sweet pepper	Promotes somatic embryogenesis.	100
	BcBBM	Chinese white poplar	Promotes somatic embryogenesis.	101
GRF	AtGRF5, HaGRF5, GmGRF5, BnGRF5, BvGRF5, ZmGRF5, ZmGRF5-LIKE1	Sugar beet, soybean, maize	Callus step required or promotes organogenesis from explants.	84
Gene combinations				
WUS BBM	ZmWUS2 ZmBBM	Maize, sorghum, rice, sugarcane, salvia, wheat	Promotes somatic embryogenesis. Use of specific maize promoters or excision required to avoid developmental defects. Uses ternary or super-binary vector.	59,67,102
GRF-GIF	TaGRF4-GIF1	Wheat, rice, triticale, Citrus, watermelon, hemp	Callus step required.	71,83,103
GRF-GIF BBM	TaGRF4-GIF1 ZmBBM	Maize	Promotes somatic embryogenesis.	89
WUS IPT WUS STM	ZmWUS2 IPT AtSTM	Tomato, potato, grape	Promotes de novo meristem initiation at wound sites. Many edited plants show developmental defects and are sterile.	66

At, *Arabidopsis thaliana*; Ta, *Triticum aestivum* (bread wheat); Zm, *Zea mays* (corn); Tc, *Theobroma cacao* (cacao); Bn, *Brassica napus* (rapeseed); Bo, *Brassica oleracea* (cabbage); Br, *Brassica rapa* (mustard); Brr, *Brassica rapa* var. *rapa* (turnip); Bc, *Brassica campestris* (mustard); Gm, *Glycine max* (soybean); Gh, *Gossypium hirsutum* (cotton); Bv, *Beta vulgaris* (sugar beet); Ha, *Helianthus annuus* (sunflower).

and WUSCHEL-RELATED HOMEBOX (WOX), which are key regulators of embryogenesis initiation and meristematic stem cell fate, as summarized in the previous section. Individual or combined expression of these regulators in somatic cells is often sufficient to trigger whole-plant regeneration under in vitro cultivation or de novo shoot formation on soil-grown plants⁶⁶. While detailed reviews on the use of morphogenic genes have been published^{3,63,65}, here we focus on the most recent advances using these strategies specifically for the transformation of crop species.

Ectopic expression of the maize *BBM* and *WUS2* genes (a co-orthologue of *Arabidopsis WUS*) increased transformation efficiency in several crops, including maize, sorghum, indica rice, sugarcane and genotypes recalcitrant to biolistic and *Agrobacterium*-mediated transformation⁵⁹. Additionally, combinatorial expression of *WUS2* together with different developmental regulators (including the cytokinin biosynthesis *IPT* and the meristematic *STM* genes) at the wound sites of soil-grown plants induced de novo meristem formation and enabled gene editing, bypassing the tissue-culture step in tomato, potato and grape⁶⁶.

However, constitutive expression of morphogenic genes is usually not well tolerated, often resulting in infertile plants or plants with undesired pleiotropic phenotypes. While these undesired effects can be segregated away for genome-editing approaches, alternative strategies to express these factors transiently (only during a specific stage) or to entirely remove them post-infection have been pursued^{3,59,67}. In a recent strategy to overcome this issue, the non-cell autonomous function of *ZmWUS2* was exploited to produce transformed plants

without the *ZmWUS2* expression cassette, a phenomenon designated as altruistic transformation. In this system, two independent strains of *Agrobacterium*, one containing the selectable marker and the other the cassette expressing *ZmWUS2*, were successfully used in different ratios for maize embryo infections⁶⁸. Similarly, a recently developed CRISPR-based approach to simultaneously induce the expression of endogenous morphogenic genes and promote editing increased the recovery of transformed and edited plants without developmental defects in poplar and rice, offering a new tool for accelerating genome engineering⁶⁹.

Recent studies have shown that constitutive expression of either *WOX5* or a GROWTH REGULATING FACTOR4 (*GRF4*) and *GRF-INTERACTING FACTOR1* (*GIF1*) chimaeric protein enhances transformation efficiency in wheat as well as several other monocot species^{70,71}. While both systems still require a callus-induction stage, they were shown to expand the range of genotypes that can be regenerated, a major limitation to the widespread and fast adoption of transformation technologies in crops. *WOX5* belongs to the same homeobox family of *WUS* and plays a key role in root stem cell maintenance, and its overexpression induces shoot regeneration in *Arabidopsis* calli, possibly by promoting *TAA1* auxin biosynthesis and interfering with cytokinin signalling⁷²⁻⁷⁶. The overexpression of a wheat *WOX5* gene significantly enhanced the transformation efficiency in recalcitrant genetic backgrounds in wheat, barley and maize, seemingly without detrimental effects on overall development, making it an attractive system for monocot transformation⁷⁰. However, in maize, *WOX5* was tested in combination with a ternary

vector system; therefore, whether *WOX5* itself is responsible for the reported efficiency has yet to be determined⁷⁷.

GRFs are highly conserved, plant-specific transcription factors that promote growth and boost organ size when overexpressed^{78–80}. GRFs form protein complexes with GIF cofactors that in turn interact with SWI/SNF chromatin remodelling factors^{81,82}. By forcing physical proximity of both proteins in the GRF–GIF chimaeric protein, the system proved efficient for transformation by increasing regeneration not only in durum and bread wheat but also in rice, triticale and the dicot crops *Citrus*, watermelon and hemp^{71,83}. Furthermore, the GRF–GIF chimaera allowed wheat shoot regeneration in media lacking cytokinin, allowing marker-free selection of transgenic plants and suggesting a potential role of the GRF–GIF complex in the regulation of endogenous cytokinins⁷¹. The use of individual GRF factors has been reported to improve transformation efficiency in sugar beet, canola, sunflower and maize without affecting the development or fertility of transgenic plants⁸⁴. While the molecular mechanisms of how GRF–GIF complexes promote regeneration are currently not fully known, the direct downstream targets of rice OsGRF6 included auxin biosynthesis and signalling genes, suggesting that GRF–GIF complexes promote regeneration by elevating auxin biosynthesis and auxin signalling in transformed cells⁸⁵. In addition, a regulatory loop between *GRF* and *PLT* genes was reported to establish the boundary between the stem cell niche and the transit-amplifying region in *Arabidopsis* root meristems, suggesting that GRF–GIF complexes could influence PLT activity⁷⁹.

Concluding remarks and future perspectives

Despite the recent breakthroughs using morphogenic factors, challenges still remain in developing fast, efficient and reliable transformation systems that can be quickly adopted by the public sector, particularly for monocot species. Given that some of these morphogenic factors function in hierarchical order during embryogenesis and meristem formation^{40,47}, their combination may promote somatic embryogenesis in an additive manner (for example, BBM WUS2)^{59,67,68,86–88} (Table 1), even when the molecular mechanisms driving these synergistic interactions are not entirely known^{59,66,67}. Indeed, preliminary data suggest that a higher efficiency of maize embryo transformation is achievable using a combination of the wheat GRF–GIF chimaera and the maize BBM transcriptional regulator with standard *Agrobacterium* binary vector systems⁸⁹ (Table 1). The identification of additional genes involved in the regeneration process may foster additional combinations that could prove optimal for the transformation of certain recalcitrant crop species or genotypes. The selection of optimal combinations would benefit from additional insights into the regeneration process, which should be obtainable from single-cell chromatin accessibility and expression data of specific species, genotypes and organs⁷⁶. These studies may also aid the identification of molecular signatures of precursor cells during somatic embryogenesis and provide new targets for biotechnological approaches to transformation.

Despite recent progress, the tissue and genotype dependency of the transformation process still represent major bottlenecks to the widespread use of transgenic and genome-editing technologies in many economically important crop species. Some of the technologies discussed here have been tested with success in different tissues, including embryos excised from dry seeds and leaf segments from seedlings³⁹. An ideal system should use reliable and easily accessible sources of tissue (for example, leaves, coleoptiles or root tips) or even protoplasts⁹⁰ for transformations, and specific morphogenic triggers to overcome their intrinsic lack of competency for somatic embryogenesis and regeneration⁴⁰. In species such as maize and wheat, this would facilitate the widespread adoption of transformation technologies in academic labs, whereas in species clonally propagated or with embryos too small to handle, it may represent the only viable strategy.

To accelerate crop transformation and breeding, it is also imperative to develop new DNA delivery methods with reduced genotype

dependency, which may involve the use of new bacterial strains with high efficiency of infection⁹¹ or the adoption of specific nanomaterials for transformation. Promising advances with ternary vector systems for *Agrobacterium*-mediated transformation^{20,21} may more broadly tackle the transformation of recalcitrant backgrounds of different crops and may be adaptable to different transformation systems. The identification of new genes and pathways that confer genotype dependency of transformation may also contribute to solving this issue, as shown in maize, where knocking out *SAUR15* (an early auxin-responsive gene) significantly increased regeneration efficiency⁴³. Pursuing a variety of these approaches will probably be key to increase the adoption of transformation technologies in recalcitrant crops and genetic backgrounds, including wild progenitors and diverse germplasms^{92,93}, for both basic and applied research.

References

- Phillips, R. L., Kaeppler, S. M. & Olhoft, P. Genetic instability of plant tissue cultures: breakdown of normal controls. *Proc. Natl Acad. Sci. USA* **91**, 5222–5226 (1994).
- Neelakandan, A. K. & Wang, K. Recent progress in the understanding of tissue culture-induced genome level changes in plants and potential applications. *Plant Cell Rep.* **31**, 597–620 (2012).
- Gordon-Kamm, B. et al. Using morphogenic genes to improve recovery and regeneration of transgenic plants. *Plants (Basel)* **8**, 38 (2019).
- Anami, S., Njuguna, E., Coussens, G., Aesaert, S. & Van Lijsebettens, M. Higher plant transformation: principles and molecular tools. *Int. J. Dev. Biol.* **57**, 483–494 (2013).
- Liu, J. et al. Genome-scale sequence disruption following biolistic transformation in rice and maize. *Plant Cell* **31**, 368–383 (2019).
- Clark, K. A. & Krysan, P. J. Chromosomal translocations are a common phenomenon in *Arabidopsis thaliana* T-DNA insertion lines. *Plant J.* **64**, 990–1001 (2010).
- Hu, Y., Chen, Z., Zhuang, C. & Huang, J. Cascade of chromosomal rearrangements caused by a heterogeneous T-DNA integration supports the double-stranded break repair model for T-DNA integration. *Plant J.* **90**, 954–965 (2017).
- Krispil, R. et al. The position and complex genomic architecture of plant T-DNA insertions revealed by 4SEE. *Int. J. Mol. Sci.* **21**, 2373 (2020).
- Pucker, B., Kleinbolting, N. & Weisshaar, B. Large scale genomic rearrangements in selected *Arabidopsis thaliana* T-DNA lines are caused by T-DNA insertion mutagenesis. *BMC Genomics* **22**, 599 (2021).
- Jupe, F. et al. The complex architecture and epigenomic impact of plant T-DNA insertions. *PLoS Genet.* **15**, e1007819 (2019).
- Woo, J. W. et al. DNA-free genome editing in plants with preassembled CRISPR–Cas9 ribonucleoproteins. *Nat. Biotechnol.* **33**, 1162–1164 (2015).
- Liang, Z. et al. Genome editing of bread wheat using biolistic delivery of CRISPR/Cas9 in vitro transcripts or ribonucleoproteins. *Nat. Protoc.* **13**, 413–430 (2018).
- Liang, Z. et al. Efficient DNA-free genome editing of bread wheat using CRISPR/Cas9 ribonucleoprotein complexes. *Nat. Commun.* **8**, 14261 (2017).
- Svitashev, S., Schwartz, C., Lenderts, B., Young, J. K. & Mark Cigan, A. Genome editing in maize directed by CRISPR–Cas9 ribonucleoprotein complexes. *Nat. Commun.* **7**, 13274 (2016).
- Hamilton, C. M., Frary, A., Lewis, C. & Tanksley, S. D. Stable transfer of intact high molecular weight DNA into plant chromosomes. *Proc. Natl Acad. Sci. USA* **93**, 9975–9979 (1996).
- De Saeger, J. et al. *Agrobacterium* strains and strain improvement: present and outlook. *Biotechnol. Adv.* **53**, 107677 (2021).

17. Lacroix, B. & Citovsky, V. Pathways of DNA transfer to plants from *Agrobacterium tumefaciens* and related bacterial species. *Annu Rev. Phytopathol.* **57**, 231–251 (2019).
18. Yuan, Z. C. et al. The plant signal salicylic acid shuts down expression of the vir regulon and activates quorum-sensing genes in *Agrobacterium*. *Proc. Natl Acad. Sci. USA* **104**, 11790–11795 (2007).
19. Lee, C. W. et al. *Agrobacterium tumefaciens* promotes tumor induction by modulating pathogen defense in *Arabidopsis thaliana*. *Plant Cell* **21**, 2948–2962 (2009).
20. Zhang, Q. et al. A novel ternary vector system united with morphogenic genes enhances CRISPR/Cas delivery in maize. *Plant Physiol.* **181**, 1441–1448 (2019).
21. Anand, A. et al. An improved ternary vector system for *Agrobacterium*-mediated rapid maize transformation. *Plant Mol. Biol.* **97**, 187–200 (2018).
22. Kang, M. et al. An improved *Agrobacterium*-mediated transformation and genome-editing method for maize inbred B104 using a ternary vector system and immature embryos. *Front Plant Sci.* **13**, 860971 (2022).
23. Raman, V. et al. *Agrobacterium* expressing a type III secretion system delivers *Pseudomonas* effectors into plant cells to enhance transformation. *Nat. Commun.* **13**, 2581 (2022).
24. Lv, Z., Jiang, R., Chen, J. & Chen, W. Nanoparticle-mediated gene transformation strategies for plant genetic engineering. *Plant J.* **104**, 880–891 (2020).
25. Vejlupkova, Z. et al. No evidence for transient transformation via pollen magnetofection in several monocot species. *Nat. Plants* **6**, 1323–1324 (2020).
26. Zhao, X. et al. Pollen magnetofection for genetic modification with magnetic nanoparticles as gene carriers. *Nat. Plants* **3**, 956–964 (2017).
27. Wang, Z. P. et al. Efficient and genotype independent maize transformation using pollen transfected by DNA-coated magnetic nanoparticles. *J. Integr. Plant Biol.* **64**, 1145–1156 (2022).
28. Ma, X., Zhang, X., Liu, H. & Li, Z. Highly efficient DNA-free plant genome editing using virally delivered CRISPR–Cas9. *Nat. Plants* **6**, 773–779 (2020).
29. Hu, J. et al. A barley stripe mosaic virus-based guide RNA delivery system for targeted mutagenesis in wheat and maize. *Mol. Plant Pathol.* **20**, 1463–1474 (2019).
30. Li, T. et al. Highly efficient heritable genome editing in wheat using an RNA virus and bypassing tissue culture. *Mol. Plant* **14**, 1787–1798 (2021).
31. Williams, L. E. Genetics of shoot meristem and shoot regeneration. *Annu. Rev. Genet.* **55**, 661–681 (2021).
32. Ikeuchi, M. et al. Molecular mechanisms of plant regeneration. *Annu. Rev. Plant Biol.* **70**, 377–406 (2019).
33. Motte, H., Vereecke, D., Geelen, D. & Werbrouck, S. The molecular path to in vitro shoot regeneration. *Biotechnol. Adv.* **32**, 107–121 (2014).
34. Efroni, I. et al. Root regeneration triggers an embryo-like sequence guided by hormonal interactions. *Cell* **165**, 1721–1733 (2016).
35. Verma, S., Attuluri, V. P. S. & Robert, H. S. An essential function for auxin in embryo development. *Cold Spring Harb. Perspect. Biol.* **13**, a039966 (2021).
36. Cheng, Y., Dai, X. & Zhao, Y. Auxin synthesized by the YUCCA flavin monooxygenases is essential for embryogenesis and leaf formation in *Arabidopsis*. *Plant Cell* **19**, 2430–2439 (2007).
37. Stepanova, A. N. et al. TAA1-mediated auxin biosynthesis is essential for hormone crosstalk and plant development. *Cell* **133**, 177–191 (2008).
38. Lardon, R., Wijnker, E., Keurentjes, J. & Geelen, D. The genetic framework of shoot regeneration in *Arabidopsis* comprises master regulators and conditional fine-tuning factors. *Commun. Biol.* **3**, 549 (2020).
39. Lin, G. et al. Chromosome-level genome assembly of a regenerable maize inbred line A188. *Genome Biol.* **22**, 175 (2021).
40. Wang, F. X. et al. Chromatin accessibility dynamics and a hierarchical transcriptional regulatory network structure for plant somatic embryogenesis. *Dev. Cell* **54**, 742–757e748 (2020).
41. Li, M. et al. Auxin biosynthesis maintains embryo identity and growth during BABY BOOM-induced somatic embryogenesis. *Plant Physiol.* **188**, 1095–1110 (2022).
42. Uc-Chuc, M. A. et al. YUCCA-mediated biosynthesis of the auxin IAA is required during the somatic embryogenic induction process in *Coffea canephora*. *Int. J. Mol. Sci.* **21**, 4751 (2020).
43. Wang, Y. et al. Genetic variations in ZmSAUR15 contribute to the formation of immature embryo-derived embryonic calluses in maize. *Plant J.* **109**, 980–991 (2021).
44. Wojcikowska, B. et al. LEAFY COTYLEDON2 (LEC2) promotes embryogenic induction in somatic tissues of *Arabidopsis*, via YUCCA-mediated auxin biosynthesis. *Planta* **238**, 425–440 (2013).
45. Lotan, T. et al. *Arabidopsis* LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells. *Cell* **93**, 1195–1205 (1998).
46. Zhang, T. Q. et al. A two-step model for de novo activation of WUSCHEL during plant shoot regeneration. *Plant Cell* **29**, 1073–1087 (2017).
47. Wu, L. Y. et al. Dynamic chromatin state profiling reveals regulatory roles of auxin and cytokinin in shoot regeneration. *Dev. Cell* **57**, 526–542e527 (2022).
48. Matsuo, N., Makino, M. & Banno, H. *Arabidopsis* ENHANCER OF SHOOT REGENERATION (ESR)1 and ESR2 regulate in vitro shoot regeneration and their expressions are differentially regulated. *Plant Sci.* **181**, 39–46 (2011).
49. Iwase, A. et al. WIND1 promotes shoot regeneration through transcriptional activation of ENHANCER OF SHOOT REGENERATION1 in *Arabidopsis*. *Plant Cell* **29**, 54–69 (2017).
50. Heyman, J. et al. The heterodimeric transcription factor complex ERF115–PAT1 grants regeneration competence. *Nat. Plants* **2**, 16165 (2016).
51. Ikeuchi, M. et al. Wounding triggers callus formation via dynamic hormonal and transcriptional changes. *Plant Physiol.* **175**, 1158–1174 (2017).
52. Sakamoto, Y. et al. Transcriptional activation of auxin biosynthesis drives developmental reprogramming of differentiated cells. *Plant Cell* **34**, 4348–4365 (2022).
53. Hofhuis, H. et al. Phyllotaxis and rhizotaxis in *Arabidopsis* are modified by three PLETHORA transcription factors. *Curr. Biol.* **23**, 956–962 (2013).
54. Kareem, A. et al. PLETHORA genes control regeneration by a two-step mechanism. *Curr. Biol.* **25**, 1017–1030 (2015).
55. Lian, Z. et al. Application of developmental regulators to improve in planta or in vitro transformation in plants. *Plant Biotechnol. J.* **20**, 1622–1635 (2022).
56. Hernandez-Coronado, M. et al. Plant glutamate receptors mediate a bet-hedging strategy between regeneration and defense. *Dev. Cell* **57**, 451–465.e6 (2022).
57. Boutilier, K. et al. Ectopic expression of BABY BOOM triggers a conversion from vegetative to embryonic growth. *Plant Cell* **14**, 1737–1749 (2002).
58. Khanday, I., Skinner, D., Yang, B., Mercier, R. & Sundaresan, V. A male-expressed rice embryogenic trigger redirected for asexual propagation through seeds. *Nature* **565**, 91–95 (2019).
59. Lowe, K. et al. Morphogenic regulators BABY BOOM and WUSCHEL improve monocot transformation. *Plant Cell* **28**, 1998–2015 (2016).

60. Khanday, I., Santos-Medellin, C. & Sundaresan, V. Rice embryogenic trigger BABY BOOM1 promotes somatic embryogenesis by upregulation of auxin biosynthesis genes. Preprint at *bioRxiv* <https://doi.org/10.1101/2020.08.24.265025> (2020).
61. Horstman, A. et al. The BABY BOOM transcription factor activates the LEC1–ABI3–FUS3–LEC2 network to induce somatic embryogenesis. *Plant Physiol.* **175**, 848–857 (2017).
62. Underwood, C. J. et al. A PARTHENOGENESIS allele from apomictic dandelion can induce egg cell division without fertilization in lettuce. *Nat. Genet.* **54**, 84–93 (2022).
63. Maren, N. A. et al. Genotype-independent plant transformation. *Hortic. Res.* **9**, uhac047 (2022).
64. Salaun, C., Lepiniec, L. & Dubreucq, B. Genetic and molecular control of somatic embryogenesis. *Plants (Basel)* **10**, 1467 (2021).
65. Kausch, A. P. et al. Edit at will: genotype independent plant transformation in the era of advanced genomics and genome editing. *Plant Sci.* **281**, 186–205 (2019).
66. Maher, M. F. et al. Plant gene editing through de novo induction of meristems. *Nat. Biotechnol.* **38**, 84–89 (2020).
67. Lowe, K. et al. Rapid genotype ‘independent’ *Zea mays* L. (maize) transformation via direct somatic embryogenesis. *In Vitro. Cell. Dev. Biol. Plant* **54**, 240–252 (2018).
68. Hoerster, G. et al. Use of non-integrating *Zm-Wus2* vectors to enhance maize transformation. *In Vitro. Cell. Dev. Biol. Plant* **56**, 265–279 (2020).
69. Pan, C. et al. Boosting plant genome editing with a versatile CRISPR–Combo system. *Nat. Plants* **8**, 513–525 (2022).
70. Wang, K. et al. The gene *TaWOX5* overcomes genotype dependency in wheat genetic transformation. *Nat. Plants* **8**, 110–117 (2022).
71. Debernardi, J. M. et al. A GRF–GIF chimeric protein improves the regeneration efficiency of transgenic plants. *Nat. Biotechnol.* **38**, 1274–1279 (2020).
72. Sarkar, A. K. et al. Conserved factors regulate signalling in *Arabidopsis thaliana* shoot and root stem cell organizers. *Nature* **446**, 811–814 (2007).
73. Ortiz-Ramirez, C. et al. Ground tissue circuitry regulates organ complexity in maize and *Setaria*. *Science* **374**, 1247–1252 (2021).
74. Forzani, C. et al. *WOX5* suppresses *CYCLIN D* activity to establish quiescence at the center of the root stem cell niche. *Curr. Biol.* **24**, 1939–1944 (2014).
75. Pi, L. et al. Organizer-derived *WOX5* signal maintains root columella stem cells through chromatin-mediated repression of *CDF4* expression. *Dev. Cell* **33**, 576–588 (2015).
76. Zhai, N. & Xu, L. Pluripotency acquisition in the middle cell layer of callus is required for organ regeneration. *Nat. Plants* **7**, 1453–1460 (2021).
77. Wang, K. et al. Author correction: the gene *TaWOX5* overcomes genotype dependency in wheat genetic transformation. *Nat. Plants* **8**, 717–720 (2022).
78. Li, S. et al. The *OsmiR396c–OsGRF4–OsGIF1* regulatory module determines grain size and yield in rice. *Plant Biotechnol. J.* **14**, 2134–2146 (2016).
79. Rodriguez, R. E. et al. MicroRNA *miR396* regulates the switch between stem cells and transit-amplifying cells in *Arabidopsis* roots. *Plant Cell* **27**, 3354–3366 (2015).
80. Liebsch, D. & Palatnik, J. F. MicroRNA *miR396*, GRF transcription factors and GIF co-regulators: a conserved plant growth regulatory module with potential for breeding and biotechnology. *Curr. Opin. Plant Biol.* **53**, 31–42 (2020).
81. Debernardi, J. M. et al. Post-transcriptional control of GRF transcription factors by microRNA *miR396* and GIF co-activator affects leaf size and longevity. *Plant J.* **79**, 413–426 (2014).
82. Luo, G. & Palmgren, M. GRF–GIF chimeras boost plant regeneration. *Trends Plant Sci.* **26**, 201–204 (2021).
83. Zhang, X. et al. Establishment of an *Agrobacterium*-mediated genetic transformation and CRISPR/Cas9-mediated targeted mutagenesis in hemp (*Cannabis sativa* L.). *Plant Biotechnol. J.* **19**, 1979–1987 (2021).
84. Kong, J. et al. Overexpression of the transcription factor *GROWTH-REGULATING FACTOR5* improves transformation of dicot and monocot species. *Front. Plant Sci.* **11**, 572319 (2020).
85. Gao, F. et al. Blocking *miR396* increases rice yield by shaping inflorescence architecture. *Nat. Plants* **2**, 15196 (2015).
86. Aesaert, S. et al. Optimized transformation and gene editing of the B104 public maize inbred by improved tissue culture and use of morphogenic regulators. *Front. Plant Sci.* **13**, 883847 (2022).
87. Masters, A. et al. *Agrobacterium*-mediated immature embryo transformation of recalcitrant maize inbred lines using morphogenic genes. *J. Vis. Exp.* <https://doi.org/10.3791/60782> (2020).
88. Mookkan, M., Nelson-Vasilchik, K., Hague, J., Zhang, Z. J. & Kausch, A. P. Selectable marker independent transformation of recalcitrant maize inbred B73 and sorghum P898012 mediated by morphogenic regulators *BABY BOOM* and *WUSCHEL2*. *Plant Cell Rep.* **36**, 1477–1491 (2017).
89. Chen, Z., Debernardi, J. M., Dubcovsky, J. & Gallavotti, A. The combination of morphogenic regulators *BABY BOOM* and *GRF–GIF* improves maize transformation efficiency. Preprint at *bioRxiv* <https://doi.org/10.1101/2022.09.02.506370> (2022).
90. Reed, K. M. & Bargmann, B. O. R. Protoplast regeneration and its use in new plant breeding technologies. *Front. Genome Ed.* **3**, 734951 (2021).
91. Cho, H. J. et al. Development of an efficient marker-free soybean transformation method using the novel bacterium *Ochrobactrum haywardense* H₁. *Plant Biotechnol. J.* **20**, 977–990 (2022).
92. Zobrist, J. D. et al. Transformation of teosinte (*Zea mays* ssp. *parviglumis*) via biolistic bombardment of seedling-derived callus tissues. *Front. Plant Sci.* **12**, 773419 (2021).
93. Hufford, M. B. et al. De novo assembly, annotation, and comparative analysis of 26 diverse maize genomes. *Science* **373**, 655–662 (2021).
94. Thakare, D., Tang, W., Hill, K. & Perry, S. E. The MADS-domain transcriptional regulator *AGAMOUS-LIKE15* promotes somatic embryo development in *Arabidopsis* and soybean. *Plant Physiol.* **146**, 1663–1672 (2008).
95. Arroyo-Herrera, A. et al. Expression of *WUSCHEL* in *Coffea canephora* causes ectopic morphogenesis and increases somatic embryogenesis. *Plant Cell Tissue Organ Cult.* **94**, 171–180 (2008).
96. Che, P. et al. *Wuschel2* enables highly efficient CRISPR/Cas-targeted genome editing during rapid de novo shoot regeneration in sorghum. *Commun. Biol.* **5**, 344 (2022).
97. Liu, Y. et al. Establishment of *Agrobacterium*-mediated genetic transformation and application of CRISPR/Cas9 genome-editing system to *Brassica rapa* var. *rapa*. *Plant Methods* **18**, 98 (2022).
98. Hu, W. et al. *Kn1* gene overexpression drastically improves genetic transformation efficiencies of citrus cultivars. *Plant Cell Tissue Organ Cult.* **125**, 81–91 (2016).
99. Elhiti, M., Tahir, M., Gulden, R. H., Khamiss, K. & Stasolla, C. Modulation of embryo-forming capacity in culture through the expression of *Brassica* genes involved in the regulation of the shoot apical meristem. *J. Exp. Bot.* **61**, 4069–4085 (2010).
100. Heidmann, I., de Lange, B., Lambalk, J., Angenent, G. C. & Boutilier, K. Efficient sweet pepper transformation mediated by the *BABY BOOM* transcription factor. *Plant Cell Rep.* **30**, 1107–1115 (2011).

101. Deng, W., Luo, K., Li, Z. & Yang, Y. A novel method for induction of plant regeneration via somatic embryogenesis. *Plant Sci.* **177**, 43–48 (2009).
102. Zhou, Z. et al. Boosting transformation in wheat by BBM–WUS. Preprint at *bioRxiv* <https://doi.org/10.1101/2022.03.13.483388> (2022).
103. Feng, Q. et al. Highly efficient, genotype-independent transformation and gene editing in watermelon (*Citrullus lanatus*) using a chimeric ClGRF4–GIF1 gene. *J. Integr. Plant Biol.* **63**, 2038–2042 (2021).

Acknowledgements

Research in the Gallavotti lab is supported by grants from the National Science Foundation (IOS nos 1546873, 1916804 and 2026561).

Research in the Dubcovsky lab is supported by grants no. 2022-68013-36439 and no. 2022-67013-36209 from the USDA National Institute of Food and Agriculture and by the Howard Hughes Medical Institute.

Author contributions

Z.C., J.M.D., J.D. and A.G. conceived the manuscript, contributed to writing and editing, and approved the manuscript.

Competing interests

J.M.D. is co-inventor in patent no. US2017/0362601A1, which describes the use of chimaeric GRF–GIF proteins with enhanced effects on plant growth (Universidad Nacional de Rosario Consejo Nacional de

Investigaciones Científicas y Técnicas). J.D. and J.M.D. are co-inventors in UC Davis patent application no. WO2021007284A2, which describes the use of GRF–GIF chimaeras to enhance regeneration efficiency in plants.

Additional information

Correspondence should be addressed to Andrea Gallavotti.

Peer review information *Nature Plants* thanks David Jackson and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2022